



# **GYNORIT** گائینورٹ

**TABLETS** گولیاں

**Progestogen-estrogen combination**

Important information, please read carefully:

## **COMPOSITION:**

1 Gynorit tablet contains 10 mg norethisterone acetate B.P. and 0.02 mg. ethinylloestradiol B.P.

## **INDICATIONS:**

Secondary amenorrhoea of short duration (up to about 1 year).

Treatment with preparations such as

Gynorit tablets is to be given at the earliest 8 weeks after the last menstrual period and after pregnancy has been excluded.

## **DOSAGE AND ADMINISTRATION:**

Pregnancy must be excluded before the administration of Gynorit tablets. One Gynorit tablet is taken whole with a little liquid on each of two consecutive days.

Gynorit tablets usually induces a menstruation like withdrawal bleeding within 3-6 days, in isolated cases later. Further diagnostic and therapeutic measures are required if no bleeding occurs within 2 weeks.

## **CONTRA-INDICATIONS:**

Pregnancy

A history of herpes of pregnancy

## **PRESENTATION:**

2 tablets

Store all drugs properly and keep them out of reach of children.

---

Manufactured by:  
LAHORE CHEMICAL & PHARMACEUTICAL  
WORKS (PVT.)LTD., (Authorised users)  
137-Ferozepur Road, Lahore.