IFLOR SACHET 250 / CAP 250 (SACCHAROMYCES BOULARDII)

ايس فلور عهدا اليول ١٥٠٠

(LYOPHILIZED)

COMPOSITION:

Each ENFLOR sachet contains: Each ENFLOR capsule contains: Saccharomyces boulardii 250mg (Lyophilized).

PROPERTIES - MECHANISM OF ACTION:

ENFLOR is a new biotherapeutic agent. It is a French research product, which acts in intestine in a living state. The pharmacological studies show that ENFLOR has multiple favorable actions e.g.,

- Microbial antagonism: In vitro and in vivo microbial antagonism against bacteria e.g. E. coli, Salmonella spp., Shigella spp., and candida albicans.
- Stimulation of Immune defenses: ENFLOR induces a significant increase in the concentration of the secretory component and IgAs (the first line of defense against infection in the digestive tract) in the small intestine and may turn out to have a protective effect against infections of digestive origin.
- Antitoxin activity: It acts on the origin of the secretion of enterotoxins and / or reduces the enterotoxic effects of toxins by inhibition toxin receptor binding sites.
- Enzymatic activity: It produces a significant increase of disaccharidases including Lactase, sucrase, and

INDICATIONS:

- Treatment of infectious and non specific diarrhea. ENFLOR can also be used in conjunction with oral re-hydration for the treatment of acute diarrhea.
- Treatment and prevention of digestive complication of antibiotic therapy: diarrhea, colitis, and
- Treatment and prevention of nasogastric-tube alimentation diarrhea.

1 to 2 sachets given once or twice daily. 1 to 2 Capsules given once or twice daily. These doses may be increased in accordance with the prescription of a physician without any contraindication.

DIRECTION FOR USE:

Sachet: Pour the contents (Powder) of the sachet into some water, beverage or semi liquid.

CAUTION:

ENFLOR should not be poured on or mixed with very cold or very hot liquid or food.

FOR CONTINUOUS ENTERAL FEEDING:

Add ENFLOR to the nutrient solution at the time of preparation. It can be stored at room temperature for several days.

INTERACTIONS:

ENFLOR should not be administered with systemic anti-fungai agents.

SIDE EFFECTS:

Some rare epigastric disturbances have been reported, not requiring that treatment be discontinued.

INSTRUCTIONS:


Store in a cool, dry place and protect from sunlight Keep away from reach of children.

PRESENTATION:

Box of 10 sachets. Box of 1x 10's strip.

ایک سےدوسافے پاکھیول روزانہ ایک سےدوم تبہ طريقة استعال ساشے: ساشے ميں موجود ياؤوركو پائى جلول يارقيق (تيكى)غذام ملائميں اور في ليں-طريقة استعال كيدول: كيدول كوياني كساته ليس-این فلورکو بہت شخندی یا گرم اشیاء (محلول یا غذا) کے ساتھ شلایں۔ خندى اورخنگ جگه بر بچول كى دسترس سے دُور ركيس اورسورج كى روشى سے بحائيں-

Manufactured by: Hilton Pharma (Pvt.) Ltd. 13, Sector 15, Korangi Industrial Area, Karachi, Pakistan.



Martin Dow